



THE BIRD CLINIC VETERINARY CORPORATION

NUTRITIONAL PROBLEMS IN PET BIRDS

Tradition dictates that pet birds eat predominately, if not exclusively, a seed diet. Sadly, this long outdated tradition has created a situation which promotes severe nutritional imbalance and medical problems in our pet and aviary birds. Old feeding traditions ("wives tales") have become so entrenched in today's society that it seems like heresy to alter these "accepted" methods of care. Any of the topics below can cause serious malnutrition, poor immunity, illness, and a much shortened lifespan (as much as 40%) for our pet birds.

Commercial formulated pelleted foods solve these issues!

THE VEGETARIAN TYPE DIET

A "classic" **seed diet is a vegetarian diet** still promoted by many people in the pet industry. Unfortunately, **seeds contain protein of low biological value**, as do grains, fruits, greens, and vegetables. The amino-acid structure of these foods is deficient in both quantity and quality of essential amino acids, making this vegetarian diet hazardous. Deficiencies of essential amino acids easily can and do occur, especially in Cockatoo species. In addition, many scientific papers have demonstrated that seed diets provide only 20% of the necessary calcium levels for an adult bird. It is not surprising that many new patients presented to The BIRD Clinic suffer from metabolic bone disease, egg binding, and other signs of hypocalcaemia. They also present with anemia and poor muscle growth from amino acid deficiencies.

THE UNBALANCED DIET

Ingesting only one food group will always lead to nutritional deficiencies. Seed diets cannot be scientifically formulated because seeds come from natural sources and only represent one food group. Birds eating these seeds with a few fruits and vegetables are at high risk of malnutrition. Some commercial bird seed companies add pellets to their mixtures in an attempt to balance out these deficiencies. Unfortunately, the mixture **ONLY** provides a balanced diet if **EVERY** part of the mix is eaten. These added pellets are not balanced or tested for texture or palatability, so it is rare that a bird will eat them. Some diets do not even list their ingredients for comparison and evaluation. Homemade customized seed diets often have been put together with the best of intentions and the best of ingredients, but trying to balance a bird's diet without training in nutrition to at least a Ph.D. level is like trying to fix a car without a manual. Diets need to be evaluated through a process of calculations, chemical analysis, and testing including testing in controlled feeding trials with LIVE birds for an extended period (3-5 years) to fully evaluate whether the analysis is indeed correct. Since the early 1990's several feeding trials have been performed by reputable companies. Dr. Nemetz personally performed a comparative nutritional study over a six-year period in a breeding colony of Meyer's Parrots and his pet Meyer's, Gigi, is celebrating 30 years of health as of this writing.

CONFINEMENT IN SMALL CAGES

An animal's nutritional status can be indirectly affected by continuous caging. An avoidance behavior pattern develops in which a bird shuns anything that is not familiar. Because of this, the bird rejects any new foods. In doing this, the bird clings to his narrow diet and is fearful of new foods which could help to balance his/her diet. This can be overcome by repeated offerings of new foods in a similar location or several locations in the bird's cage over a two-week or longer period of time. This allows the bird an adjustment period to investigate the "new" cage item and has proven very successful in the majority (>98%) of cases.

You simply need to be very patient and do not give up! You will win over time.....



THE BIRD CLINIC VETERINARY CORPORATION

FREE CHOICE FEEDING

Placing a variety of foods in a dish and letting birds eat free choice naturally leads to problems. Birds do not have the instinct to pick a balanced diet from a selection of foods. When observing the feeding behavior of pet birds one will notice they have dietary "binges". That is, **they will eat one food almost exclusively for a period of time and then without obvious reason switch to another food and eat it in preference to anything else**. Partially because of inherited characteristics, and partially from acquired habits, birds can become locked into eating a narrow selection of foods. They will approach their food dish as if more concerned about what not to eat rather than what they should eat. **Picking and pecking, spilling out, sorting through, and finally selecting a food, makes up their normal native daily foraging routine.** This highly undesirable situation fosters nutritional deficiencies in addition to wasting more food than they eat. Unfortunately, these poor eating habits may go unrecognized when birds are allowed to eat free choice from a variety of foods constantly in front of them. Whether from being a "picky" eater or from *monophagism* (habitual eating of one kind of food), a bird eating mainly one food (except a balanced pelleted food) receives too much of some nutrients and too little of others, causing body cells to suffer and even die. At The BIRD Clinic it has been shown over the years that **we can achieve 40%+ longer life expectancy** from our pet birds if we have them eat a good quality, studied, formulated pelleted diet.

FAST METABOLIC RATE

Although pet and aviary birds spend many hours in cages, their metabolism and structure are that of a flying machine. A healthy bird burns a hot fire internally with an average core temperature of 105°F; manufactures more heat than any other vertebrate; consumes more oxygen; eats more food per unit body weight; and moves more quickly. All of these factors confirm the unusually fast metabolism of a bird. Because of this fast metabolic rate, birds need to be fed a different diet than other animals. They require relatively large volumes of calorically dense, highly digestible food which constitutes a nutritionally complete diet. **Their instinctive need for energy is one reason birds will search out high calorie products like seeds and nuts, which are unfortunately not highly digestible or nutritionally complete.** Compounding this problem is that the energy requirement of a caged pet bird is much less than their wild counterparts. **Parrots in the wild spend half of each and every day foraging** for food and water, expending a great deal of energy. Further, because of the large "turn over" in food daily (with food passing from mouth to feces in approximately 3-5 hours); what may be a minor deficiency in other animals become magnified into a major deficiency in a bird. Try thinking of the bird as a super-athlete requiring a super-diet!

CONCLUSION

Many inaccurate, if not false, ideas still exist about the proper nutrition of pet birds. We still do not have all the answers, as there are over 50 species of birds common in the pet bird industry, but now with **over 50 years of research**, observation and testing, great strides have been made and will continue through time. **Health-wise, pet birds are at an advantage in eating a completely balanced, commercially formulated, pelleted diet 90% by volume of their total diet, with the remaining 10% (by volume) as healthy human food treats** – proteins, complex carbohydrates, vegetables, and minimal fruits. Table food is an adjunct to a well-balanced diet. It adds interest, flavor, and freshness to a balanced commercially pelleted diet, but must only be given in very small daily quantities.

*** CAUTION / WARNING ***

**Never abruptly remove any existing diet for a new diet assuming the bird will convert.
Birds can and have died from starvation using this method.**

*This publication is part of The BIRD C News™ series to help educate bird owners. Produced by The BIRD Clinic Veterinary Corporation
200 S. Tustin Street, Suite E, Orange, CA 92866-2322 (714) 633-2910. www.thebirdclinic.com May be reproduced with permission only.*